

**13.1 Exploring Periodic Data**


**Periodic function** - repeats a pattern of y-values (outputs) at regular intervals.

**Cycle** - 1 complete pattern. A cycle may begin at any point on the graph of the function.

**Period** - the horizontal length of 1 cycle, - in terms of x-values.

**Example 1: Identifying Cycles and Periods**


Analyze the periodic function. a) Identify 1 cycle in two different ways. b) Determine the period of the function.



$1 - (-3) = 4 \text{ units}$        $3 - 0 = 3 \text{ units}$

**Example 2: Identifying Periodic Functions**

Determine whether each function is or is not periodic. If it is, find the period.




*not periodic*

*periodic*  
*9 units*

$9 - 0$

**Amplitude** - of a periodic function measures the amount of variation in the y-values. To find the amplitude:

**Example 3: Find the Amplitude of the periodic function.**




$$\text{amp: } \frac{\text{max} - \text{min}}{2} = \frac{1 - (-2)}{2} = \frac{3}{2}$$

$$\text{amp: } \frac{\text{max}_y - \text{min}_y}{2} = \frac{3 - 0}{2} = \frac{3}{2}$$

13.2 Angles (day 1)

An angle in standard position has:

- vertex is at the origin
- one ray is on the x-axis



**Initial side** - the ray on the positive x-axis


**Terminal side** - the other ray of the angle

The **measure of an angle in standard position** is the amount of rotation from the initial side to the terminal side.

The measure of an angle is **positive** when the rotation from the initial side to terminal side is counterclockwise


The measure of an angle is **negative** when the rotation from the initial side to terminal side is clockwise

Sketch an angle in standard position:



**Example 1: Measuring an Angle in Standard Position**


Find the measure of the angle.




**Example 2: Sketching an Angle in Standard Position**

Sketch each angle in standard position.


a.  $85^\circ$



b.  $-320^\circ$



c.  $180^\circ$



**Coterminal Angles** - Two angles in standard position are coterminal if they have the same terminal side. For any given angle, there are infinitely many coterminal angles. To find a coterminal angle, add or subtract  $360^\circ$ .

**Example 3: Finding Coterminal Angles**

a. Find a positive angle and a negative angle that are coterminal with  $198^\circ$ .

$$198^\circ + 360^\circ = 558^\circ$$

$$198^\circ - 360^\circ = -162^\circ$$

b. Are the angles with measure  $40^\circ$  and  $680^\circ$  coterminal? Explain.

$40^\circ$   $680^\circ$  not coterminal. the difference is not a multiple of  $360^\circ$ .

$$680 - 40 = \frac{640}{360} = X$$


c. Find the measure of an angle between  $0^\circ$  and  $360^\circ$  coterminal with  $385^\circ$ .

$$385^\circ - 360^\circ = 25^\circ$$

d. Find the measure of an angle between  $0^\circ$  and  $360^\circ$  coterminal with  $-356^\circ$ .


$$-356^\circ + 360 = 4^\circ$$

**Recall: Special Right Triangles**




Find the missing side lengths in each  $45^\circ-45^\circ-90^\circ$  triangle. Rationalize any denominators.


1. hypotenuse = 1 inch



2. leg = 2 cm




3. hypotenuse =  $\sqrt{3}$  ft




Find the missing side lengths in each  $30^\circ-60^\circ-90^\circ$  triangle. Rationalize any denominators.


4. shorter leg = 3 inch



5. longer leg = 1 cm



6. hypotenuse = 1 ft




$\theta$  angle


13.2 The Unit Circle (day 2)

Warm Up: Sketch the angle in standard position:


1.  $\theta = 30^\circ$



2.  $\theta = 45^\circ$



3.  $\theta = 60^\circ$





Unit Circle - a circle with a radius of 1 and its center is at the origin

Points on the unit circle are related to periodic functions. You can use the symbol  $\theta$  "theta" for the measure of an angle in standard position.

Sketch the unit circle:

$$\frac{\text{adj}}{\text{hyp}} = \frac{\sqrt{3}}{2}$$

$$\frac{\text{opp}}{\text{hyp}} = \frac{1}{2}$$



$$\cos \theta = \left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$$

$$\sin \theta = \frac{1}{2}$$

$$\sin 30^\circ = \frac{\text{opp}}{\text{hyp}} = \frac{1}{2}$$

SOHCAHTOA


Definition: Cosine and Sine of an Angle

Given: an angle in standard position with measure  $\theta$ .

The cosine of  $\theta$  is the  $x$ -coordinate of the point at which the terminal side intersects the unit circle.

The sine of  $\theta$  is the  $y$ -coordinate of the point at which the terminal side intersects the unit circle.

Sketch:




$$\cos 60^\circ = \frac{1}{2}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

Example 4&5: Find the cosine and sine of the angle. Provide a sketch of the angle in standard position.


a.  $30^\circ$




$$\cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\sin 30^\circ = \frac{1}{2}$$

b.  $45^\circ$



c.  $60^\circ$



$$\frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{\sqrt{2}}$$


$$\frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{\sqrt{2}}$$

$$\sin = \frac{\sqrt{2}}{2} = \frac{\sqrt{2}}{2}$$


$$\cos 45^\circ = \frac{\sqrt{2}}{2} = \frac{\sqrt{2}}{2}$$

Find the cosine and sine of the angle. Provide a sketch of the angle in standard position.


d.  $0^\circ$




e.  $90^\circ$




f.  $120^\circ$



g.  $-120^\circ$



h.  $-60^\circ$



i.  $150^\circ$

**Practice Problems:**

Calculator Needed: For angles that are not a multiple of  $30^\circ$  or  $45^\circ$ , you will need your calculator. Find  $\cos \theta$  and  $\sin \theta$ .

1.  $\theta = 32^\circ$

$\cos 32^\circ = 0.834$   
 $\sin 32^\circ = 0.551$

2.  $\theta = -210^\circ$

$\cos(-210^\circ) = -0.884$   
 $\sin(-210^\circ) = -0.468$

3.  $\theta = -10^\circ$

Find a positive and negative coterminal angle for the given angle.  $\pm 360^\circ$


4.  $\theta = 400^\circ$

$(+360) = 760^\circ$   
 $400 - 360 = 40^\circ$   
 $40 - 360 = -320^\circ$

5.  $\theta = -125^\circ$


$-125 + 360 = 235^\circ$   
 $-125 - 360 = -485^\circ$

6.  $\theta = -57^\circ$




In which quadrant, or on which axis, does the terminal side of each angle lie? Sketch the angle to help you.


7.  $210^\circ$




8.  $-60^\circ$




9.  $270^\circ$



h.  $-60^\circ$ 

$$\sin(-60^\circ) = -\frac{\sqrt{3}}{2}$$

$$\cos(-60^\circ) = \frac{1}{2}$$

i.  $150^\circ$ 


$$y: \sin 150^\circ = \frac{1}{2}$$

$$x: \cos 150^\circ = -\frac{\sqrt{3}}{2}$$

$(\cos \theta, \sin \theta)$

# Fill in The Unit Circle

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{y}{x}$$





13.3 Radian Measure (Day 1)

Warm Up: Find the circumference of a circle with the given radius or diameter. Round your answer to the nearest tenth.

1. radius 4 in

$$C = 2\pi r$$

$$C = 2\pi \cdot 4 = 8\pi$$

$$25.1 \text{ in}$$

2. diameter 70 m


$$C = d\pi = 2\pi r$$

$$C = 70\pi$$

$$219.9 \text{ m}$$

Central angle - an angle whose vertex is the center of a circle



sketch:



Intercepted arc - the portion of the circle with endpoints on the sides of the central angle and remaining points within the interior of the angle.

Radian - when the intercepted arc equals the radius, the measure of the angle is 1 radian


sketch:



- The circumference of a circle is  $2\pi r$ . Thus there are  $2\pi$  radians in any circle.
- Since  $\frac{2\pi}{2}$  radians =  $\frac{360}{2}$ °, then  $\pi$  radians =  $180$ °.
- Thus you can use this proportion to convert between degrees and radians.

$$D \rightarrow R: \frac{\pi}{180}^\circ$$

$$R \rightarrow D: \frac{180^\circ}{\pi}$$



Example 1: Use a proportion

a. Find the radian measure of 60°.

$$60^\circ \cdot \frac{\pi}{180^\circ} = \frac{\pi}{3}$$

b. Find the degree measure of  $\frac{5\pi}{2}$  radians

$$\frac{5\pi}{2} \cdot \frac{180^\circ}{\pi} = 450^\circ$$

**Converting between Radians and Degrees**

- To convert degrees to radians, multiply by  $\frac{\pi}{180^\circ}$
- To convert radians to degrees, multiply by  $\frac{180^\circ}{\pi}$

**Example 2: Using Dimensional Analysis**

Convert the angle to degrees. Round to the nearest degree.

a.  $-\frac{3\pi}{4}$  radians

$$-\frac{3\pi}{4} \cdot \frac{180^\circ}{\pi} = -135^\circ$$

b.  $\frac{\pi}{2}$  radians

$$\frac{\pi}{2} \cdot \frac{180^\circ}{\pi} = 90^\circ$$

c. 2 radians

$$2 \text{ radians} \cdot \frac{180^\circ}{\pi \text{ rad.}}$$

$$\frac{360^\circ}{\pi}$$

Convert the angle to radians. Round to the nearest hundredth.

d.  $27^\circ$

$$27^\circ \cdot \frac{\pi}{180^\circ} = \frac{3\pi}{20} \text{ radians}$$

e.  $225^\circ$

$$225^\circ \cdot \frac{\pi}{180^\circ} = \frac{5\pi}{4} \text{ radians}$$

f.  $150^\circ$

$$150^\circ \cdot \frac{\pi}{180^\circ} = \frac{5\pi}{6} \text{ radians}$$

**Example 3: Find the exact values of  $\cos\theta$  and  $\sin\theta$  for each angle measure.**

Step 1: Convert to degrees.


Step 2: Draw the angle. The terminal side is the hypotenuse.

Step 3: Complete the right triangle. Draw a leg to the  $x$ -axis.

Step 4: State the  $\cos\theta$  and  $\sin\theta$ .


a.  $\frac{\pi}{4}$  radians

$$\frac{\pi}{4} \cdot \frac{180^\circ}{\pi} = 45^\circ \rightarrow \frac{\pi}{4}$$




b.  $\frac{\pi}{6}$  radians

$$\frac{\pi}{6} \cdot \frac{180^\circ}{\pi} = 30^\circ \rightarrow \frac{\pi}{6}$$




c.  $\frac{\pi}{2}$  radians

$$\frac{\pi}{2} \cdot \frac{180^\circ}{\pi} = 90^\circ \rightarrow \frac{\pi}{2}$$



d.  $\frac{5\pi}{6}$  radians

$$\frac{5\pi}{6} \cdot \frac{180^\circ}{\pi} = 150^\circ$$



**13.3 Arc Length (Day 2)**


**Warm Up:** Convert the angle to degrees. Draw the angle in standard position, as well as the corresponding right triangle. Then find the exact values for  $\cos\theta$  and  $\sin\theta$ .

$\theta = \frac{\pi}{3}$

You can find the length of an intercepted arc by using the proportion:

Unit Circle  $r = 1$

$S = \theta$



$C = 2\pi r$

$C = 2\pi$

$C = 2 \cdot \pi \cdot 2 = 4\pi$

$C = 2 \cdot \pi \cdot 5 = 10\pi$

**Length of an Intercepted Arc**

For a circle of radius  $r$  and a central angle of measure  $\theta$  (in radians), the length  $s$  of the intercepted arc is:


$S = \theta r$

**Example 4: Finding the Length of an Arc**


Find the length of the intercepted arc to the nearest tenth. Sketch a diagram!

a. Given: A circle of radius 3 in,  $\theta = \frac{5\pi}{6}$ .

b. Given: A circle of radius 5m,  $\theta = \frac{2\pi}{3}$ .




$s = \theta r$   
 $s = \frac{5\pi}{6} (3)$   
 $s = 7.9 \text{ in}$




$s = \theta \cdot r$   
 $s = \frac{2\pi}{3} \cdot 5$   
 $\frac{10\pi}{3} \text{ m}$   
 $10.5 \text{ m}$

The Unit Circle: radius = 1

If you know quadrant 1, you can derive quadrants 2, 3, 4 by symmetry. Thus, let's study quadrant 1.



Now let's do all four quadrants...




**13.4 The Sine Function (Day 1)**

Warm Up: Use the graph. State:

1. the period  $360^\circ$  or  $2\pi$
2. the domain  $\mathbb{R}$
3. the amplitude  $1$
4. the range  $-1 \leq y \leq 1$   $[-1, 1]$

sine function  $y = \sin\theta$ : for each measure of  $\theta$ , the sine of  $\theta$  corresponds with the  $y$ -coordinates on the unit circle.

$y = \sin\theta$	$0$	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\pi$	$\frac{3\pi}{2}$	$2\pi$
$\theta$ (degrees)	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$90^\circ$	$180^\circ$	$270^\circ$	$360^\circ$
$y$	$0$	$\frac{1}{2}$	$\frac{\sqrt{2}}{2} \approx 0.707$	$\frac{\sqrt{3}}{2} \approx 0.866$	$1$	$0$	$-1$	$0$




**Example 1: Interpreting the Sine Function in Degrees**

- a. What is the value of  $y = \sin\theta$  for  $\theta = 270^\circ$ ?  $-1$
- b. For what values of  $\theta$  between  $0^\circ$  and  $360^\circ$  does the graph of  $y = \sin\theta$  reach
  - o the maximum value of  $y = 1$ ?  $90^\circ$
  - o the minimum value of  $y = -1$ ?  $270^\circ$
  - o x-intercept of  $y = 0$ ? aka "zero"  $0^\circ, 180^\circ, 360^\circ$

**Mathematical convention:** An angle measure  $\theta$  can be expressed in degrees or in radians. However, when no unit is mentioned, you should use radians. Let's graph the sine function in radians...

$y = \sin \theta$

$\theta$ (radians)					
y					



**Example 2: Estimating Sine Values in Radians**

Use your graph above to estimate the value. Check your estimate with a calculator.

a.  $\sin 2$

0.8  
0.909

b.  $\sin \pi$

0

For the sine function, find the following:

a. amplitude

1

b. period (in degrees and radians)

$360^\circ$   
 $2\pi$

c. domain and range

D:  $\mathbb{R}$   
R:  $[-1, 1]$   
from to  
 $[0, 5)$   
 $0 \leq y < 5$

$-1 \leq y \leq 1$

**Properties of Sine Functions**

Suppose  $y = a \sin b\theta$ , where  $a \neq 0$ ,  $b > 0$ , and  $\theta$  in radians.

- The amplitude of the function is  $a$
- The number of cycles in the interval from 0 to  $2\pi$  is  $b$
- The period of the function is  $\frac{2\pi}{b}$   $\frac{360}{b}$


$y = \sin \theta$  period  $2\pi$   
 $\theta \rightarrow 360$   
 $\sin 360$   
 $\sin 180$   
 $\sin 2\theta$  period  $\pi$

**Examples 3&4: Finding the Period and Amplitude of a Sine Function**


- Find the amplitude.
- How many cycles does the sine function have in the interval from 0 to  $2\pi$ ?
- Find the period.




The  $\theta$ -axis represents values from 0 to  $2\pi$ . Each interval on the y-axis represents 1 unit.




$a: 1$   
 $b: 1$   
 $p: \frac{2\pi}{1} = 2\pi$




$a: 1$   
 $b: 3$   
 $p: \frac{2\pi}{3}$



$a: 1$   
 $b: 4$   
 $p: \frac{2\pi}{4} = \frac{\pi}{2}$



$a: 4$   
 $b: 1$   
 $p: 2\pi$



$a: 3$   
 $b: 2$   
 $p: \frac{2\pi}{2} = \pi$



13.4 The Sine Function (Day 2)

II. Graphing Sine Functions

You can use 5 points equally spaced through one cycle to sketch a sine curve. For  $a > 0$ , this 5-point pattern is zero-max-zero-min-zero.



$$y = a \sin b\theta$$

Example 5: Sketching a Graph

Sketch one cycle of each sine curve. Then write an equation for each graph.

a. amplitude 2, period  $4\pi$ ,  $a > 0$


$$y = 2 \sin \frac{1}{2}\theta$$



$(1,0)$   
 $\cos 0 = 1$

b. amplitude 3, period 4,  $a > 0$

$$y = 3 \sin \frac{1}{2}\pi \theta$$



$$p = \frac{2\pi}{b}$$

$$4\pi = \frac{2\pi}{b}$$

$$\frac{bp}{p} = \frac{2\pi}{p}$$


$$\frac{4\pi b}{4\pi} = \frac{2\pi}{4\pi}$$

$$b = \frac{2\pi}{p}$$

$$b = \frac{1}{2}$$

$$b = \frac{2\pi}{p} = \frac{2\pi}{4} = \frac{1}{2}\pi$$

c. Predict the 5-point pattern for the sine function when  $a < 0$ . Then sketch amplitude 2, period  $2\pi/3$ .



$$y = -2 \sin 3\pi \theta$$

$$b = \frac{2\pi}{(\frac{2\pi}{3})} \cdot \frac{3}{2\pi} = 3$$

**Example 6: Graphing from a Function Rule**


Sketch one cycle of the following sine functions.

1.  $y = \frac{1}{2}\sin 2\theta$   $b = 2$

amplitude:  $\frac{1}{2}$

period:  $\pi$   $p = \frac{2\pi}{b} = \frac{2\pi}{2}$

interval spacing on  $\theta$ -axis:  $\frac{\pi}{8}$



5-point pattern:




2.  $y = 3\sin \frac{\pi}{2}\theta$

amplitude:

period:

interval spacing on  $\theta$ -axis:

5-point pattern:




3.  $y = -4\sin \frac{1}{2}\theta$

amplitude:

period:


interval spacing on  $\theta$ -axis:

5-point pattern:



**13.5 The Cosine Function (Day 1)**

**Warm up:** Fill out the unit circle. Evaluate the following angles of cosine.





- 1.  $\cos 0^\circ$
- 2.  $\cos 90^\circ$
- 3.  $\cos 180^\circ$
- 4.  $\cos 270^\circ$
- 5.  $\cos 360^\circ$
- 6.  $\cos 0$
- 7.  $\cos \frac{\pi}{2}$
- 8.  $\cos \pi$
- 9.  $\cos \frac{3\pi}{2}$
- 10.  $\cos 2\pi$

Remember:  $\cos\theta$  equals the \_\_\_\_\_-coordinate on the unit circle.

**I. Graph the cosine function:  $y=\cos\theta$**

$\theta$ (radians)					
y					



**Example 1: Interpreting the Graph of  $y=\cos\theta$**

a. Use your graph above. Find the following:

- domain
- period
- range
- amplitude

b. In the interval from 0 to  $2\pi$ , where do the maximum occur? minimum? zeros?

c. What is the 5-point pattern of the cosine graph?

**Properties of Cosine Functions**

Suppose  $y = a \cos b\theta$ , where  $a \neq 0$ ,  $b > 0$ , and  $\theta$  in radians.

- The amplitude of the function is \_\_\_\_\_
- The number of cycles in the interval from 0 to  $2\pi$  is \_\_\_\_\_
- The period of the function is \_\_\_\_\_

**Example 2: Sketching the Graph of a Cosine Function**

Sketch one cycle of the following cosine functions.


1.  $y = \cos \frac{\pi}{2} \theta$

amplitude:

period:

interval spacing  
on  $\theta$ -axis:

5-point pattern:




2.  $y = -3 \cos \theta$

amplitude:

period:

interval spacing  
on  $\theta$ -axis:

5-point pattern:




3.  $y = 1.5\cos 2\theta$

amplitude:

period:

interval spacing  
on  $\theta$ -axis:

5-point pattern:



**13.5 The Cosine Function (Day 2)**

Warm up: Graph the following cosine functions.


1.  $y = 3\cos 2\theta$

amplitude:

period:

interval spacing  
on  $\theta$ -axis:

5-point pattern:




2.  $y = -2\cos \theta$

amplitude:

period:

interval spacing  
on  $\theta$ -axis:

5-point pattern:

**II. Solving Trigonometric Equations****Example 4: Solving a Cosine Equation**

Solve the cosine equation in the interval from 0 to  $2\pi$ . Round to the nearest hundredth. Calculators needed.

a.  $-2\cos \theta = 1.2$

b.  $3\cos 2t = -2$

c.  $5\cos \frac{7}{2}t = 3$

Identify the period, range, and amplitude of each function.

22.  $y = 3\cos\theta$

24.  $y = 2\cos\frac{1}{2}t$

26.  $y = 3\cos\left(-\frac{\theta}{3}\right)$

28.  $y = 16\cos\frac{3\pi}{2}t$

**13.6 The Tangent Function**

**Warm up:** Use a calculator to find the sine and cosine of each  $\theta$ . Then calculate the ratio of  $\sin\theta$  to  $\cos\theta$ .


$\theta$	$\sin\theta$	$\cos\theta$	$\frac{\sin\theta}{\cos\theta}$
1. $\frac{\pi}{3}$			
2. $30^\circ$			
3. $90^\circ$			
4. $\pi$			
5. $\frac{7\pi}{6}$			

**I. The Tangent Function**

The  $\cos\theta$  is derived from the \_\_\_\_\_ - coordinate of the point on the unit circle.


The  $\sin\theta$  is derived from the \_\_\_\_\_ - coordinate of the point on the unit circle.

The  $\tan\theta$  is derived from the ratio of  $\sin\theta$  to  $\cos\theta$ . In other words:  $\tan\theta = \underline{\hspace{2cm}}$



**$y=\tan\theta$**

$\theta$ (radians)	0	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	$\pi$	$-\frac{\pi}{4}$	$-\frac{\pi}{2}$
y							



**Features of the parent tangent function:**

passes through:

each "branch" is:

x-intercepts (zeros):

vertical asymptotes:

"guide points":



**Properties of Tangent Functions**

Suppose  $y = a \tan b\theta$ , where  $b > 0$ , and  $\theta$  in radians.

- The period of the function is \_\_\_\_\_
- 1 cycle occurs in the interval from \_\_\_\_\_ to \_\_\_\_\_
- There are vertical asymptotes at each end of the cycle.
- The pattern is "asymptote, -a, zero, a, asymptote".

**Example 2: Graphing a Tangent Function**

Sketch 2 cycles of each tangent function.

1.  $y = \tan \pi \theta$


period:

1 cycle: from \_\_\_\_\_ to \_\_\_\_\_

VA:

2 guide points:

pattern:



2.  $y = \tan 3\theta$


period:

1 cycle: from \_\_\_\_\_ to \_\_\_\_\_

VA:

2 guide points:

pattern:



3.  $y = \tan \frac{\pi}{2} \theta$


period:

1 cycle: from \_\_\_\_\_ to \_\_\_\_\_

VA:

2 guide points:


pattern:



**13.7 Translating Sine and Cosine Functions**

**Warm Up:**

1. Graph  $y = \tan \theta$  ... again! (Try not to peek at prior notes.)



**Features of the tangent function:**

passes through:

each "branch" is:

x-intercepts (zeros):

vertical asymptotes:

"guide points":

2. Compare each pair of equations. State the translations (horizontal, vertical) involved.

a.  $y = 2x$ ,  $y = 2x + 5$

b.  $y = |x|$ ,  $y = |x + 3|$

c.  $y = x^2$ ,  $y = x^2 - 4$

d.  $y = |x - 2| + 1$

e.  $y = x^2$ ,  $y = (x + 3)^2 - 6$

d.  $y = f(x)$ ,  $y = f(x - h) + k$

**I. Graphing Translations of Trigonometric Functions**

**Phase shift** - the horizontal translation of a function. If  $f(x)$  is the "parent", then  $f(x-h)$  translates horizontally  $h$  units. For example:  $f(x-1)$  translates \_\_\_\_\_,  $f(x+3)$  translates \_\_\_\_\_.

**Vertical shift** - the vertical translation of a function. If  $f(x)$  is the "parent", then  $f(x)+k$  translates vertically  $k$  units. For example:  $f(x) - 1$  translates \_\_\_\_\_,  $f(x) + 3$  translates \_\_\_\_\_.

**Example 1: Identifying Phase Shifts and Vertical Shifts**

What is the value of  $h$  and  $k$  in each translation? Describe the shift i.e. "3 units to the left".

a.  $f(x-2)$

b.  $y = \cos(x+4)$

c.  $f(t-5)$

d.  $y = \sin(x+3)$

e.  $f(x) - 2$

b.  $y = \cos x + 4$

c.  $f(t) - 5$

d.  $y = \sin x + 3$

**Example 2: Graphing Translations**


Make a table and then graph the following functions on the same set of axes:

$y = \sin x$

x					
y					

$y = \sin x + 3$

x					
y					




$y = \cos x$

x					
y					

$y = \cos(x - \frac{\pi}{2})$

x					
y					



**Example 3: Graphing a Combined Translation**

1.  $y = \sin(x + \pi) - 2$

amplitude:


period:

phase shift:

interval spacing on x-axis:

vertical shift:

5 point pattern:



2.  $y = 2\cos(x - \frac{\pi}{2}) + 3$

amplitude:


period:

phase shift:

interval spacing on x-axis:

vertical shift:

5 point pattern:



**Summary: Families of Sine and Cosine Functions**

Parent	Transformed Function
--------	----------------------

$y = \sin x$	_____
--------------	-------

$y = \cos x$	_____
--------------	-------

amplitude =	h =
-------------	-----

period =	k =
----------	-----

**13.7 Translating Sine and Cosine Functions (Day 2)**

**Warm Up:**

$$y = -3\sin\left(x + \frac{\pi}{2}\right) + 2$$

amplitude:


period:

phase shift:

interval spacing on x-axis:

vertical shift:

5 point pattern:



**When the phase shift is a pesky number...**

$$y = \sin\left(x - \frac{\pi}{3}\right)$$

amplitude:


period:

phase shift:

interval spacing on x-axis:

vertical shift:

5 point pattern:



**Graphing a translation of  $y = \sin 2x$ ...**

$$y = \sin 2\left(x - \frac{\pi}{3}\right) - \frac{3}{2}$$

amplitude:


period:

phase shift:

interval spacing on x-axis:

vertical shift:

5 point pattern:



$$y = -3\sin 2\left(x - \frac{\pi}{3}\right) - \frac{3}{2}$$

amplitude:


period:

phase shift:

interval spacing on x-axis:

vertical shift:

5 point pattern:



$$y = 2\cos\frac{\pi}{2}(x + 1) - 3$$


amplitude:

period:

phase shift:

interval spacing on x-axis:

vertical shift:



5 point pattern:

**Example 5: Writing a Translation**

Write an equation for each translation.

a.  $y = \sin x$ ,  $\pi$  units down

b.  $y = -\cos x$ , 2 units left

c.  $y = \cos x$ ,  $\frac{\pi}{2}$  units up

d.  $y = 2\sin x$ ,  $\frac{\pi}{4}$  units right



**13.8 Reciprocal Trigonometric Functions****Warm up:**

Find the reciprocal of each fraction:

1.  $\frac{9}{13}$

2.  $-\frac{5}{8}$

3.  $\frac{1}{2\pi}$

4.  $\frac{14}{-7}$

5.  $\theta$

Name the 3 trigonometric functions you have studied so far:

1. \_\_\_\_\_

2. \_\_\_\_\_

3. \_\_\_\_\_

These 3 trigonometric functions have reciprocals.

**Definition: Cosecant, Secant, and Cotangent****Example 1: Using Reciprocals**

a. Use your calculator (degree mode). Round your answer to the nearest hundredth.

$\csc 60^\circ$

$\cot 55^\circ$

$\sec 15^\circ$

b. Suppose  $\cos\theta = \frac{5}{13}$ . Find  $\sec\theta$ .c. Suppose  $\sin\theta = \frac{-12}{13}$ . Find  $\csc\theta$ .**Example 2: Find The Exact Value**

$\csc 30^\circ$

$\csc 45^\circ$

$\csc 60^\circ$

$\csc 90^\circ$

$\sec 30^\circ$

$\sec 45^\circ$

$\sec 60^\circ$

$\sec 90^\circ$

$\cot 30^\circ$

$\cot 45^\circ$

$\cot 60^\circ$

$\cot 90^\circ$

**Example 3: Using Radians**

a. Use your calculator (radian mode). Round your answer to the nearest hundredth.

$$\sec(-1)$$

$$\csc(-1.5)$$

$$\sec 2$$

b. Find the exact value.


$$\cot \frac{\pi}{3}$$

$$\cot \pi$$

$$\sec 0$$

**Example 4: Graph The Reciprocal Trigonometric Functions**

<b>x</b>	0	$\frac{\pi}{2}$	$\pi$	$\frac{3\pi}{2}$	$2\pi$
<b>y = sin x</b>					
<b>y = csc x</b>					



<b>x</b>	0	$\frac{\pi}{2}$	$\pi$	$\frac{3\pi}{2}$	$2\pi$
<b>y = cos x</b>					
<b>y = sec x</b>					

